

in Urbanized Eastern

Massachusetts 02601 USA; e-mail: jw9802@

eastern Massachusetts. Canadian Field-

in heavily urbanized eastern Massa-
100.5 km and averaged 63.8 ± 52.0 km
Transients ranged in age between 1-2 yr
($r = 0.389$, $P = 0.518$). Coyotes travel long
distances. Because of the ability of Coyotes to
move on educating the public about actual

movements, suburban, transient, urban-

those regions; conversely, if the op-
Coyotes move similar distances in
areas), then control efforts would likely
unless targeting a specific individ-
part of a larger ongoing study on
Way et al. 2001; Way et al. 2002a;
in eastern Massachusetts, documents
transient Coyotes in a heavily urban-

took place in two urbanized locations:
the towns and cities north of Boston
research conducted on the heavily
of Boston (~ 100 - 150 km²; 42.43° N,
in the cities of Revere (3089 peo-
density = $1318/\text{km}^2$), Everett (4345
ing density = $1817/\text{km}^2$), and Malden
housing density = $1800/\text{km}^2$) (U.S.
2000 estimates). The area is charac-
terized by high-density housing with small woodland
(cemeteries) non-strategically situated
areas. Coyotes were captured and spent
time in these wooded, green areas as the
open areas were often fenced and pro-
hibited Coyotes to travel, except for main
roads and holes in some of the fences
corridors between some of the green
areas (Way et al. 2006).

Research was conducted within Barnstable
County, Massachusetts (approximate study
area with a concentration in the town/city of
Barnstable, a town. Barnstable is techni-
cally 41.70° N, 70.28° W; land area = 155.5 km²).

Human population density in the town/ city of Barnstable was 308 people/km² and housing density was $161/\text{km}^2$, while the entire Barnstable County (3382 km²) averaged 217 people/km² and 144 houses/km² (U.S. Census Bureau 2000 estimates). The town/ city of Barnstable has a distinct rural-urban gradient within its borders; the highest and lowest densities of people were found in urban Hyannis (556 people/km², housing units = $328/\text{km}^2$) and rural West Barnstable ($89/\text{km}^2$, housing units = $39/\text{km}^2$) (Cape Cod Commission 1998*). Road density, defined as centerline km of roadway per km², was 4.7 for the town of Barnstable and 4.0 for Barnstable County (Cape Cod Commission 1998*). Cape Cod is characterized by being residential as well as having numerous small (5 - 10 ha) and a few large (~ 1000 ha) conservation areas interspersed throughout. Most of the neighborhoods are not fenced, however, and Coyotes were readily able to travel through these areas to access various portions of their home range (Way et al. 2004). Coyote pack territories were roughly 30 km² and were non-overlapping, similar to more rural areas (Gese et al. 1996; Way et al. 2002a).

Methods

Coyotes were captured by box trap (Way et al. 2002b) then radio-collared or radio-implanted (juveniles – i.e., pups of the year) using Telonics, Inc. (Mesa, Arizona) transmitters, aged based on tooth wear (Bowen 1982; Landon et al. 1998), weighed, blood drawn (ca. 4 cc), then released. Transient (or nomadic) Coyotes were classified as Coyotes who had no discernable territory and nomadically moved throughout the study areas, including within resident collared Coyote home ranges. These Coyotes are typically classified as young Coyotes that are in the process of dispersing from their natal pack (Way et al. 2002a).

Tracking protocols were described by Way et al. (2002a) and Way et al. (2004). Portable receivers (Custom Electronics, Urbana, Illinois, USA) and hand-held 3-element Yagi antennas were used to radio-track Coyotes both on foot and from a vehicle. Due to the highly developed landscape with many roads, I mostly radio-tracked in a vehicle as Coyotes did not react negatively to them as much as they did to people (e.g., by running away; Way 2007a; J. Way, unpublished data); occasionally I approached radio-collared Coyotes as closely as possible on foot without disturbing them. I used binoculars and video-cameras when observing Coyotes, and city street lights, nightscopes and occasionally headlights when following Coyotes at night with a vehicle (Way et al. 2002a; Way et al. 2004). Due to funding constraints, I did not use airplanes to search for Coyotes that left our study areas; extended trips were made in vehicles to locate missing Coyotes but this was often unsuccessful. Instead, I relied on recovering Coyotes from sightings by the public ($n = 1$), opportunistically receiving radio-locations in

new areas ($n = 1$), and recovering carcasses from human-related kills (i.e., road-kill or gunshot; $n = 3$). Transient movement distances were calculated from where they were first captured (if not on natal territory) or from their natal territories (if known) to their final location when they either settled (i.e., established a resident home range in a new area) or died.

I used an independent sample two-tailed t-test to detect differences between male and female dispersal. Levene's test was used to detect for equal variance between samples; a significant Levene's result indicated that equal variances were not assumed. I correlated Coyote dispersal distances with body weight using two-tailed bivariate Pearson Correlation Coefficients (SPSS Inc., Chicago, Illinois) tests. I set significance at < 0.05 .

Results and Discussion

I documented the movement of five transient Coyotes: one from north Boston and four from Cape Cod (Figure 2). Upon release, all Coyotes appeared to be in the process of dispersal as their movements were nomadic. Movements from capture to final location varied from 23.0 to 100.5 km and averaged 63.8 ± 52.0 km for two females and 38.7 ± 17.2 km for three males (Levene's Test = 0.030 , $t = 0.657$, $df = 1.15$, $P = 0.618$). Although I had a low sample size, these movements were within the range of dispersal by transient Coyotes in more rural environments, with an average distance of 98 km for Coyotes in forested Maine (Harrison 1992), 16 – 152 km in rural Ontario (Kolenosky et al. 1978), 20 – 140 km in rural, agricultural Vermont (Person 1988), ≤ 30 km (depending on age class) in farmland-forested central Alberta (but up to 100 – 150 km; Nellis and Keith 1976), 12.7 – 17.9 km in the western United States (Robinson and Grand 1958), 35.7 km in Iowa (Andrews and Boggess 1978), 16 – 68 km (average = 48 km) in northern Minnesota (Berg and Chesness 1978), 36.4 km in New Mexico (Young and Jackson 1951), and 40.3 – 45.6 km (up to 161 km) in Wyoming (Young and Jackson 1951).

There was no difference between male and female transient movements and all were young (1 – 2 yr old) animals (Table 1), which is typical of when canids disperse (Harrison 1992; Mech and Boitani 2003a). Although I had a low sample size, it is noteworthy that a female exhibited the longest movements. Coyote #BN0402 traveled through nearly all of eastern Massachusetts, one of the most densely human populated areas in the country (U. S. Census Bureau 2000 estimates; Way 2007a). Additionally, female Coyote #0202 navigated across a 1 km canal to leave Cape Cod (Way 2002). However, caution should be taken with these low sample sizes and over-interpreting the results. For instance, as this paper went to press, Way (2008*) documented a yearling male Coyote that travelled 81.1 km, from the village of Centerville to




FIGURE 1. Map of the two Coyote study sites in eastern Massachusetts with polygons representing all of the towns in Massachusetts.

Hingham, Massachusetts, about 15 km southeast of Boston. This one animal's movement would have caused male Coyotes in this study to go from an average dispersal distance of 38.7 to 49.3 km. Thus, the male/female difference is not as important as the fact that this study is the first to document transient Coyote movements in urbanized locals.

Harrison (1992) found no sex specific statistical differences in Coyote dispersal in his Maine study, which might be expected for a monogamous species, but the maximal distance traveled (342 km) was by a female. Other studies corroborate these findings with individual females dispersing the farthest in their studies, such as 17.9 (vs. 12.7) km in the western United States (Robinson and Grand 1958), 323.2 km in Iowa

(Andrews and Boggess 1978), 154 km in central Alberta (Nellis and Keith 1976), and 544 km from Riding Mountain National Park, Manitoba, to Saskatchewan, the furthest movement of a coyote on record (Carbyn and Paquet 1986). In fact, all long-distance movements were made by females except a male Coyote in Rosatte's (2002) study that moved 320 km in an agricultural region of southern Ontario (the most urban of the study sites after mine). However, most researchers claimed that these long-distance dispersals were rare.

The Boston Coyote likely exhibited a relatively straight-line dispersal through Boston and south to the edge of the Atlantic Ocean in southeastern Massachusetts where either the ocean stopped her movement and/or she paired up with a mate (Figure 2; Way

TABLE 1. Movements of transient (or nomadic) Coyotes in eastern Massachusetts

Coyote ID	Sex/Age (yr)	Start of dispersal ¹	Conclusion of dispersal ²	Body weight (kg)	Minimum distance traveled (km)
#BN0402	Female (1)	April 2004	November 2004	13.6	100.5
#0301	Male (2)	December 2003	February 2005	18.5	57.1
#0401	Male (2)	February 2004	December 2005	14.5	23.0
#0202	Female (2)	January 2002	May 2002	19.1	27.0
#0205	Male (1.5)	March 2002	June 2003	15.2	36.1

¹Represents when the animal was documented as a transient Coyote in the act of dispersal. Because all animals in this study were captured as transients and were likely off their natal ranges when captured, their respective capture location notes where I recorded their start of dispersal.

²Represents when an animal terminated nomadic movements or was recovered dead.

2007a). Because possible for Coyotes and because I out of radio comment direction, were documented Cod, radiating area (Figure 1) ending up both site (Figure 2). that "In homogeneity probably dispersal no habitat type Wolf density, a doubt play vary




FIGURE 2. Movements of transient (or nomadic) Coyotes in eastern Massachusetts.

2007a). Because of the restricted movement paths possible for Coyotes dispersing on peninsular Cape Cod and because I did not sequentially track them when out of radio contact, it was difficult to assess movement direction. However, the four combined Coyotes were documented to travel throughout most of Cape Cod, radiating from the town of Barnstable study area (Figure 1), with two Coyotes each ultimately ending up both east and west of the Barnstable study site (Figure 2). Mech and Boitani (2003a: 15) noted that "In homogeneous habitat types, Wolves would probably disperse equally in all directions. However, no habitat type is homogeneous, and topography, Wolf density, and areas of human development no doubt play varying roles in steering dispersal, direc-

tion." Harrison (1992) noted that Coyotes in Maine often follow major rivers and that those water barriers often deflected and influenced Coyote dispersal, while Gese and Mech (1991) found that Wolves in Minnesota dispersed in relatively equal directions.

There was no relationship between Coyote body weight and distance traveled ($r = 0.389$, $P = 0.518$) with two relatively heavy and three light transient Coyotes (see Way 2007b for weight ranges), although caution should again be taken when inferring results because of low sample size. However, Gese and Mech (1991) also found no difference between wolf body weight and the age at dispersal or dispersal success. Yet Gese et al. (1996) discovered that Coyotes that dispersed from their natal pack in Yellowstone National

Minimum distance traveled (km)
100.5
57.1
23.0
27.0
36.1

Animals in this study location notes

Park were often low ranking and had less access to food. One would presume that these Coyotes also weighted less than philopatric Coyotes.

Distances traveled by nomadic Coyotes in eastern Massachusetts represented minimum values because (1) most movements were made in a restricted, linear landscape (i.e., Cape Cod); (2) it was not known where all of the transients began dispersal (i.e., we did not capture them as juveniles within their natal ranges); and (3) it was not known if all localized (i.e., established or joined a pack) before death. If they were killed before settling then their potential dispersal distance may have been underestimated. Based on the time (months to one to two years) that I was not in contact with radio-collared Coyotes, I suspect that many Coyotes traveled the length of Cape Cod and only infrequently came into my study area. Coyotes can move long distances in short periods of time (Way et al. 2004) and I sporadically located these transients before they disappeared, presumably out of my study area.

The most accurate way to determine Coyote dispersal and subsequent transient movements is to radio-tag juveniles in their natal area and track how Coyotes disperse (e.g., Harrison 1992). However, dispersal is time intensive, expensive, and one of the most difficult of all population processes to document (Harrison 1992; Mech and Boitani 2003b). For example, I also radio-tagged nine additional juveniles (seven from Cape Cod; two from north of Boston) within their natal ranges but recovered no dispersal information on any of them because they either died on their natal territory ($n = 5$) or disappeared and were never re-located ($n = 4$). Thus, reporting transient Coyote movements in a range of landscapes, regardless of the sample size, is important.

In highly developed regions (e.g., eastern Massachusetts), it is likely that Coyotes travel across a gradient of urbanization (from rural to urban) when dispersing. In other words, an "urban Coyote" would merely represent where that individual currently lives; that same individual could have originated from a much more rural location. For example, Coyote #BN0402 was captured in the process of dispersal in heavily urban north Boston (Revere, Massachusetts). She traveled southwest into heavily urban Cambridge, then vanished before being discovered in Dartmouth, Massachusetts (Way 2007a), a town that is rural-suburban with a large percentage of open space remaining (population density = 192/km², housing density = 71 km²; U.S. Census Bureau 2000 estimates). She paired up with a male Coyote and produced pups in Dartmouth before dying of unknown causes in June 2005 in her recently established range.

Coyotes have colonized much of North America, including urbanized areas, south of the Arctic. While there may be times when individual(s) Coyotes may need to be removed from a population, widespread

control efforts have never been successful at regulating Coyote numbers (Parker 1995), partly because of their well-documented dispersal and movement ability in a myriad of landscapes. The Coyote's persistence is nearly guaranteed throughout its range and the potential for (re)colonization of habitats ranging from rural to urban is very high. Therefore, I recommend that Coyote management efforts focus more on educating the public about actual Coyote behavior (e.g., Way 2005) and their life history needs than on killing them. As noted by Gompper (2002), the presence of Coyotes in northeastern North America and in cities, represents a natural range expansion of the species, not an exotic invasion, and the species is here to stay.

Acknowledgments

This study would not have been possible without the help from Tara Way, the Way family, Eric Strauss, and the Hyannis Animal Hospital (specifically, Larry Venezia and Paul McCartin) for the Cape Cod component of the research; the Saugus Animal Hospital (specifically, Bob Binder), David Eatough and Stephen Cifuni and his family for the Boston component of our study; and Boston College's Environmental Studies Program and Urban Ecology Institute for equipment purchases. Permits and protocols have been approved and obtained through Boston College and the Massachusetts Division of Fisheries and Wildlife to handle the animals, and from numerous private, town, and state lands that allowed us to put traps in the field and to track collared Coyotes. I appreciate all of these entities that provided documents and support, especially the towns/cities of Barnstable (including rangers, administrators, and animal control), Melrose, Revere, Everett, Malden, and Saugus, among others, for allowing me to use all or some of their land as if it were my own. I thank Gary Clabaugh for reporting the sighting of #BN0402 in his backyard in Dartmouth, Massachusetts, and for civic employees for reporting the carcasses of three others.

Documents Cited (marked * in text)

- Cape Cod Commission.** 1998. Cape trends: demographic and economic characteristics and trends. Cape Cod Commission, Barnstable, Massachusetts, USA.
- Way, J. G.** 2008. Field Update January 13, 2008 - Amazing news: Coyote Walnut is 50 miles away from Centerville (where he was born) in the town of Hingham about 10 miles south-southeast of Boston. <http://easterncoyotersearch.com/update2008-01-13.html>. Retrieved on 17 July 2008.

Literature Cited

- Andrews, R. D., and E. K. Boggess.** 1978 (Reprint 2001). Ecology of Coyotes in Iowa. Pages 249-265 in *Coyotes: Biology, Behavior, and Management*. Edited by Marc Bekoff. Blackburn Press, Caldwell, New Jersey, USA.
- Berg, W. E., and R. A. Chesness.** 1978 (Reprint 2001). Ecology of Coyotes in northern Minnesota. Pages 229-247 in *Coyotes: Biology, Behavior, and Management*. Edited by Marc Bekoff. Blackburn Press, Caldwell, New Jersey, USA.

Bowen, W. *latrans*,
dian Fie
Carbyn, L.
ment of
Journal
Chubbs, T.
eastern
Naturali
Fritts, S. E.
Journal
Gese, E. M.
(*Canis*
Canadia
Gese, E. M.
and nu
dent co
Gompper
logical
northea
185-19
Harrison
coyote
128-13
Kolenosh
Wolve
ural R
Landon,
Mech,
for gra
682.
Mech, L.
Pages
Edited
go Pr
Mech, L.
behav
go Pr
Nellis, C.
Coyo
Mana
Parker,
Nimb
page
Person.
food
regio
mon

- successful at regulating (e.g., partly because of movement ability). Coyote's persistence in range and the potential for ranging from rural to urban areas. I recommend that more on educating behavior (e.g., Way on killing them, the presence of Coyotes in cities, reputation of the species, not is here to stay.
- en possible without family, Eric Strauss, (specifically, Larry the Cape Cod community Animal Hospital atough and Stephen ston component of environmental Stud-Institute for equip-ols have been ap-on College and the ies and Wildlife to erous private, town, ut traps in the field reciate all of these and support, espe-(including rangers,), Melrose, Revere, g others, for allow-land as if it were for reporting the ard in Dartmouth, oyes for reporting
- (x)
- trends: demographic nds. Cape Cod Com-USA.
- 13, 2008 - Amazing way from Centerville ngham about 10 miles sterncoyotersearch. ved on 17 July 2008.
- 1978 (Reprint 2001), 249-265 in *Coyotes: nt. Edited by Marc New Jersey, USA.*
- (Reprint 2001). Ecol-ta. Pages 229-247 in nagement. Edited by ell, New Jersey, USA.
- Bowen, W. D.** 1982. Determining age of coyotes, *Canis latrans*, by tooth sections and tooth-wear patterns. *Canadian Field-Naturalist* 96: 339-341.
- Carbyn, L. N., and P. C. Paquet.** 1986. Long distance movement of a coyote from Riding Mountain National Park. *Journal of Wildlife Management* 50: 89.
- Chubbs, T. E., and F. R. Phillips.** 2002. First record of an eastern coyote, *Canis latrans*, in Labrador. *Canadian Field-Naturalist* 116: 127-129.
- Fritts, S. H.** 1983. Record dispersal by a wolf from Minnesota. *Journal of Mammalogy* 64: 166-167.
- Gese, E. M., and L. D. Mech.** 1991. Dispersal of wolves (*Canis lupus*) in northeastern Minnesota, 1969-1989. *Canadian Journal of Zoology* 69: 2946-2955.
- Gese, E. M., R. L. Ruff, and R. L. Crabtree.** 1996. Social and nutritional factors influencing the dispersal of resident coyotes. *Animal Behavior* 52: 1025-1043.
- Gompper, M. E.** 2002. Top carnivores in the suburbs? Ecological and conservation issues raised by colonization of northeastern North America by coyotes. *BioScience* 52: 185-190.
- Harrison, D. J.** 1992. Dispersal characteristics of juvenile coyotes in Maine. *Journal of Wildlife Management* 56: 128-138.
- Kolenosky, G. B., D. R. Voigt, and R. O. Standfield.** 1978. Wolves and Coyotes in Ontario. Ontario Ministry of Natural Resources publication, Ontario, Canada. 18 pages.
- Landon, D. B., C. A. Waite, R. O. Peterson, and L. D. Mech.** 1998. Evaluation of age determination techniques for gray wolves. *Journal of Wildlife Management* 62: 674-682.
- Mech, L. D., and L. Boitani.** 2003a. Wolf social ecology. Pages 1-34 in *Wolves: behavior, ecology, and conservation. Edited by L. D. Mech and L. Boitani.* University of Chicago Press, Chicago, Illinois, USA.
- Mech, L. D., and L. Boitani.** Editors. 2003b. *Wolves: behavior, ecology, and conservation.* University of Chicago Press, Chicago, Illinois, USA.
- Nellis, C. H., and L. B. Keith.** 1976. Population dynamics of Coyotes in central Alberta, 1964-1968. *Journal of Wildlife Management* 40: 389-399.
- Parker, G. R.** 1995. Eastern coyote: the story of its success. Nimbus Publishing, Halifax, Nova Scotia, Canada. 254 pages.
- Person, D. K.** 1988. Home range, activity, habitat use, and food habits of eastern coyotes in the Champlain valley region of Vermont. University of Vermont, Burlington, Vermont, USA.
- Robinson, W. B., and E. F. Grand.** 1958. Comparative movements of bobcats and coyotes as disclosed by tagging. *Journal of Wildlife Management* 22: 117-122.
- Rosatte, R. C.** 2002. Long distance movement by a coyote, *Canis latrans*, and red fox, *Vulpes vulpes*, in Ontario: Implications for disease-spread. *Canadian Field-Naturalist* 116: 129-131.
- Thomas, H. H., and R. L. Dibblee.** 1986. A coyote, *Canis latrans*, on Prince Edward Island. *Canadian Field-Naturalist* 100: 565-567.
- Way, J. G.** 2002. Radio-collared coyote crosses Cape Cod Canal. *Northeast Wildlife* 57: 63-65.
- Way, J. G.** 2005. Assessing student learning and interest in eastern coyotes. Ph.D. dissertation, Boston College, Massachusetts, USA.
- Way, J. G.** 2007a. Suburban howls: Tracking the eastern Coyote in Urban Massachusetts. Dog Ear Publishing, Indianapolis, Indiana, USA. 338 pages.
- Way, J. G.** 2007b. A comparison of body mass of *Canis latrans* (Coyotes) between eastern and western North America. *Northeastern Naturalist* 14(1): 111-124.
- Way, J. G., P. J. Auger, I. M. Ortega, and E. G. Strauss.** 2001. Eastern coyote denning behavior in an anthropogenic environment. *Northeast Wildlife* 56: 18-30.
- Way, J. G., and D. L. Eatough.** 2006. Use of "micro"-corridors by Eastern Coyotes (*Canis latrans* var.) in a heavily urbanized area: implications for ecosystem management. *Canadian Field-Naturalist* 120(4): 474-476.
- Way, J. G., I. M. Ortega, and P. J. Auger.** 2002a. Eastern coyote home range, territoriality and sociality on urbanized Cape Cod, Massachusetts. *Northeast Wildlife* 57: 1-18.
- Way, J. G., I. M. Ortega, P. J. Auger, and E. G. Strauss.** 2002b. Box-trapping eastern coyotes in southeastern Massachusetts. *Wildlife Society Bulletin* 30: 695-702.
- Way, J. G., I. M. Ortega, and E. G. Strauss.** 2004. Movement and activity patterns of eastern coyotes in a coastal, suburban environment. *Northeastern Naturalist* 11: 237-254.
- Young, S. P., and H. H. T. Jackson.** 1951. *The Clever Coyote.* Stackpole Company, Harrisburg, Pennsylvania, USA and Wildlife Management Institute, Washington, D.C., USA. 411 pages.

Received 17 April 2007

Accepted 25 July 2008